

**INFORMED CONSENT PROCEDURES: RESPONSIBILITIES OF  
RESEARCHERS IN DEVELOPING COUNTRIES<sup>1</sup>**

**Soledad Sánchez<sup>1</sup>, Gloria Salazar<sup>2</sup>, Marcia Tijero<sup>3</sup>, and  
Soledad Díaz<sup>4</sup>**

**Instituto Chileno de Medicina Reproductiva (ICMER)**

**José Ramón Gutiérrez 295, Depto 3**

**Phone: 56-2-632 1988 / 638 4086**

**Fax: 56-2-633 6204**

**E-mail: icmer@terra.cl**

**Santiago de Chile**

**Paper presented at the Fifth International Congress of  
Bioethics London, September 2000, in the session Research  
in the Third World**

---

<sup>1</sup> Informed consent procedures: Responsibilities of researchers in developing countries. S. Sánchez, G. Salazar, M. Tijero and S. Díaz. *Bioethics* 15:398-412, 2001.

## **INFORMED CONSENT PROCEDURES: RESPONSIBILITIES OF RESEARCHERS IN DEVELOPING COUNTRIES**

### **ABSTRACT**

We describe the informed consent procedures in a research clinic in Santiago, Chile, and a qualitative study that evaluated these procedures. The recruitment process involves information, counseling and screening of volunteers, and three or four visits to the clinic. The study explored the decision-making process of women participating in contraceptive trials through thirty-six interviews.

Women understood research as experimentation or progress. The decision to participate was facilitated by the information provided; time to consider it and to discuss it with partners or relatives; and perceived benefits such as quality of care, non-cost provision of methods and medical care. For some women, participation was an opportunity to express altruism. The main obstacles for participation were perceived side effects or risks. The final risk-benefit balance was strongly influenced by women's needs.

Women perceived that the consent form benefited the clinic, proving that participants had made a free decision, and benefited the volunteers by warranting their right to free medical care. The most important problem detected was occasional misunderstanding of the information given in the form.

We concluded that a full decision-making process enhances women's ability to exercise their right to choose, and assures research institutions that trials are conducted without coercion and that the participants are committed to the study. Researchers have the responsibility of conducting this process.

## **INTRODUCTION**

The rights of participants in clinical research are a matter of concern for many actors at the international level, particularly when research is conducted in developing countries where the participants are usually vulnerable subjects, with little awareness of their rights and many times illiterate. These societies are hierarchical and people find it difficult to resist authorities. In addition, the paternalistic biomedical model implies a power imbalance between researchers and subjects. Many times, external agents seeking their own interests fund the research. Research

projects include informed consent documents that are usually prepared in a developed country language and they should be literally translated and applied to subjects of a different cultural background (1). Because of these concerns, international agencies have produced ethical guidelines to orient the researchers on the ethical aspects involved (2-4).

Researchers are the inter-phase between the project and the participants. They are responsible of selecting research issues that are relevant to the country and/or the population under study. They also should conduct the research process with respect to participant's rights, ensuring informed consent.

In this paper Informed Consent (IC), is defined as the process by which a person, having information and choices, freely accepts or refuses to take part in a clinical trial. She/he has the right to deliberation, and respect for her/his values and preferences. This process involves intellectual and affective dimensions. Participants should clearly understand the objectives and procedures of the study, their rights and commitments, and the benefits and risks involved (2-5). In addition to providing technical information, the IC process should include the

discussion of the needs and goals of a potential participant, as well as the opportunity to discuss with partners, or other relevant persons.

However, IC is frequently understood as the result of this process: the act of reading and signing a consent form. As a consequence, more attention has been paid to the content and use of the form than to the process that supports free decisions. Sometimes the IC documents proposed by foreign agencies are difficult to understand to women in developing countries. For this reason, some researchers consider them a barrier to recruitment of subjects, rather than an instrument to ensure their rights.

This paper describes the development of the informed consent procedures at the research clinic of the Instituto Chileno de Medicina Reproductiva (ICMER) in Santiago, Chile, and its evaluation through a qualitative study. This evaluation was focused on the opinions, attitudes and needs regarding the IC process among women who participated in clinical trials for the development or evaluation of contraceptive methods (6). The evaluation was designed (SS) and conducted (GS and MT) by social scientists not involved in the clinic work<sup>2</sup>.

---

<sup>2</sup> SS, psychologist; GS, psychologist, formerly with a women's health NGO; MT, sociologist, formerly with a human rights NGO.

## **THE INFORMED CONSENT PROCEDURES AT ICMER'S CLINIC**

The clinic conducts research and provides services to women and children from a low socio-economic level. Volunteers are usually urban literate women looking for contraceptive methods or other reproductive health services. They come because quality of care is not always good in public clinics (7) and effective contraceptives are much needed in a country where abortion is illegal.

The group started thirty years ago, before there was open international discussion of the bio-ethical issues related to this research. The motivation to undertake contraceptive development was to contribute to women's health and choices. Midwives selected for their caring attitudes were recruited as the staff of the clinic.

The development of the informed consent procedures has been a learning experience for the researchers and the clinic staff. Several factors contributed to consider these procedures as important, such as respect for other people and personal freedom. "When we started, the rule was do to others what you want to be done, do not what you do not want to be done. Some mistakes derived from this rule: Personally (SD<sup>3</sup>), I like to be informed, so I overloaded women with information that

---

<sup>3</sup> SD, former director of ICMER's clinic

they could not process, they had nowhere to “anchor” because they knew little about their own bodies. It took us time to realize that the particular needs and interests of each woman were different.” In addition, the staff was so convinced of the importance and benefits of the research, that they might have been sometimes too enthusiastic emphasizing its good aspects.

On the other hand, the staff has made efforts to ensure the rights and well being of the participants. Since the start, they considered volunteers as co-protagonists of the research project, in the sense that both women and staff have a role to fulfill in the research process. Therefore, they have tried to make women feel important, considered and well treated. This is reflected in the perception that the clients have of the clinic and some have said, “Here, I am treated as a person” (7).

The researchers have learned from the experience, from the women themselves, from the clinic staff that has conveyed women’s concerns, and from women’s health activists who have contributed with different perspectives. They have had guidance from an Ethical Committee created when research was initiated, and now formed by twenty persons with different backgrounds (biomedical and psychosocial scientists, women’s health advocates, and representatives of the community).

They have also been guided by international agencies sponsoring the research and international guidelines for medical research were applied when they became available.

Over the years, the recruitment process of the volunteers has become increasingly complex, demanding more tests and visits to the clinic before the actual initiation of a contraceptive method.

*Phases of the IC Process:*

Potential volunteers come spontaneously to ICMER's clinic, usually referred by someone who has participated in a previous trial. Women's first contact with the clinic is a group session about human reproduction and contraceptive methods (including all available in the country and where). This broader information given so that women could better understand the facts related to the research methods and procedures. ICMER's objectives are explained, and characteristics of ongoing clinical studies are presented. After this session, each woman has an individual interview with a midwife to explore her reproductive goals and medical history, and to give her detailed information about her initial method choice. Methods available free of charge correspond to clinical trials<sup>4</sup>.

---

<sup>4</sup> Contraceptives available at the clinic and health care are offered to women who do not meet the trial requirements or do not want to participate, for a fee similar to that of



At a second visit, women have an individual counseling session about all aspects of the trial of their interest and they have the opportunity to express their doubts. They read the consent form, and take away a copy. A screening based on their biomedical history is performed to assess if they are eligible for the study.

At a third visit, the gynecologist verifies that the potential participant understood the objectives and procedures of the study, provides additional counseling, reviews the woman's biomedical background and performs a physical examination. At this visit, women sign the informed consent document<sup>5</sup>. Pre-admission laboratory tests are done during this visit. If the screening criteria are fulfilled, selected women enter a study, usually on their fourth visit, when they initiate the contraceptive method.

The follow-up schedule varies between specific protocols and methods, but includes at least visits at the end of the first month and every three or six months thereafter. Women are encouraged to attend the clinic at any time if they have doubts or problems. They may discontinue the trial at any time or change the contraceptive chosen, and they usually continue as clients of the clinic with a different method. However,

---

the public sector clinics. Women who want methods not available are provided with temporary contraceptive protection and referred to suitable services.

<sup>5</sup> The funding agencies consider it necessary that women read and sign the consent form before going through the screening process and clinical examination.

continuation rates in the original method chosen are usually above 95% at the end of the first year.

## **EVALUATION OF THE IC PROCEDURES**

### ***Methods***

A qualitative descriptive study was carried out, through individual semi-structured interviews with potential and actual volunteers at ICMER's clinic. This methodology allowed exploring women's ideas, feelings and attitudes regarding the IC process, and aided in the comprehension of subjects' experiences and the meanings they attribute to it (8). The aim was to interview five different women at the various steps of the IC process described, after six or more months in the trial, and after voluntarily discontinuation of participation.

### *Selection of participants:*

Potential and actual volunteers coming to the clinic between November 1997 and March 1998 were invited to participate. Recruitment was ongoing for only one clinical trial to assess the potential regulatory effect of mifepristone on the bleeding irregularities associated to Norplant<sup>R</sup> implants use. Thirty-five women interviewed were involved in this trial

and only one had participated in another. Women self-selected themselves for the interview according to their willingness to participate and their available time. Their mean age was 23 years (range 18 to 40), and their mean parity was two children (range 1 to 4). They were urban, literate and of low or middle socio-economic class. Thirty-three were in stable union, two were separated and one was single. Twenty-two were housewives, seven had a full-time job and seven had a part-time job.

*Data collection and analysis:*

Twenty interviews took place before admission (five for each phase described), five at method initiation, three after deciding not to participate in the trial, six while participating, and two after discontinuation (only two women asked for implant removal). The objectives of the interview were explained to each volunteer, who read and signed a consent form. A sociologist (MT) and a psychologist (GS) who had no previous contact with the participants conducted the interviews. Thirty-four women were interviewed during a regular visit to the clinic, in a private room, and two who had chosen not to participate in the trial were interviewed at home. Each interview lasted 20 to 30 minutes.

Interviews were taped and transcribed to word processor files. Anonymity was assured assigning a code to each transcription. The information was codified using the Ethnograph program, and classified using the categories previously defined (i.e., women's reasons to participate in clinical studies), as well as the new categories identified in subjects' speech (6). Later, an interpretative analysis was carried out, generating a picture of women's experiences during the decision-making process.

## **Results**

### *Initial Motivations:*

Women came to the clinic looking for a highly effective, long-term contraceptive. Most were referred by relatives or friends, and were motivated by cost-free contraceptive methods and the fact that ICIMER is a specialized institution:

*“My mother had used it (implants) so I more or less know what to expect, and this makes it easier for you to participate in a trial. Because at first you are afraid, but since my mother had used it and explained to me how it was...”*

Participating in a clinical trial had no special meaning at this point, besides the fact that the method they wanted was associated to an ongoing investigation.

*Processing the information received:*

The participants highly valued being informed about physiology, contraceptive methods, their side effects (topics usually avoided in formal education), and the study procedures. In general, they evaluated the information received as complete, clear, and useful.

*“The midwife explained everything to us, she explained all contraceptive methods, with great detail, and the doubts regarding Norplant that I had.”*

*“She explained it very clearly... Then you can really make a better decision, which method to use...”*

*“They even tell us we can get bruises when they take a blood sample, this is, they warn us about everything.”*

Women processed this information according to their personal needs, cultural level and cognitive abilities, as well as the information previously obtained from different sources. Women selected relevant information that made sense to them, thus receiving and processing the counseling in their own terms. Some women only paid attention to the method they were interested in. Sometimes, mistaken beliefs were not modified by the new information received, especially when they came from a significant source such as mother and friends.

Other determinants of information processing were the complexity of the contents, the timing of the information giving, and the staff's ability to convey clear data. Method's efficacy was relatively difficult for women to understand and, since it was a fundamental concern, some kept asking until satisfied. The least understood topics were method's mechanism of action and the objectives of the trial. Side effects, study requirements and visits schedule, were usually well understood.

The individual interview with a midwife was particularly important for the volunteers, because they could talk about their needs, something they often did not want to share in the group session.

Most women discussed the possibility of participating in the trial with their partner, or planned to do so. They usually read the consent form together, and some partners attended the counseling sessions. Men were involved in the decision-making process to various degrees: joint decision-making, support of women's decision, and non-participation (women decided alone, either informing them or not). Men who got involved in the process usually supported the decision of entering the trial, although some women mentioned that their partner was concerned about eventual health risks.

For most women, new doubts appeared after their first visits to the clinic. They appreciated the possibility of explaining them, and the readiness of the staff to answer the questions and clarify the doubts.

Participants considered that oral communication and written information complemented each other. The last allowed re-reading and new considerations, either alone or with relevant others, something that was appreciated.

*"I read it [the IC form] alone. Then I took it home and we both read it, he read it. At first, it looked difficult, it made me scared. But then, you begin to trust, and in the end, (you say) OK! The thing is that your own decision has a great influence, and then*

*you start seeing it in a different way, because the paper doesn't change, but the way you see it changes."*

*Perceptions of Research:*

Women associated research to experiments and progress. This had no particular affective meaning for some, while it was a strong motivation for others. Some interviewees said that participating in the clinical trial made them feel "*different*", or "*unique*". Some valued the opportunity to contribute to all women's health, or "*to a better future*", and expressed the wish to do something for others. This helped to overcome fears and concerns:

*"At first one is a little scared, but one has to try, this is science and one has to test the contraceptives, then others will know that they are good."*

On the other hand, a woman initially interested "*because research is good*" decided not to participate because she would become "*a guinea pig*," although this expression bared no negative meaning to other women. One woman's husband opposed to her participation, arguing that she would have health problems, and that experiments should be



done only with animals. Since she was not afraid, she decided to participate anyway.

*Risk-benefit balance:*

Beside contraceptive methods available, women perceived as benefits some components of the trial such as cost-free follow up and exams, money for transportation, calls from the clinic staff, and consultation on holidays. Their commitment to attend to follow-up visits was perceived as a benefit since it implied care for their own health, often left aside or seen as the last family priority.

*“I think that [being a volunteer in a trial] is good, because they take all these tests, they take good care of you. On the contrary, when you go to the public clinics they give you a contraceptive method, and you may not go back for a follow-up visit, and nobody cares.”*

Women considered high quality of care an important benefit. Friendliness and respect, ample and personalized information, and short waiting time had not been experienced by many. This gave them confidence towards the research staff and the institution, and

predisposed them to receive information about any trial with an open mind.

Most women mentioned fear to side effects and risks that they perceived associated with the method or the trial, such as weight increase, method failure, and malformed embryos in case of pregnancy. Sometimes these fears were conveyed as the opinion of the partner rather than the own woman's concern. Nevertheless, those for whom the benefits outweighed the risks were still interested in the trial.

In some cases, fears were handled during counseling, and/or by women's perception of ICMER as a specialized institution, assuming rigorous care and clinical procedures:

*"I feel safer, because this institution is dedicated to family planning, and that's a great reassurance you have. In a new clinic for instance, it's not the same ... I trust the specialists here."*

Other women accepted some risk as part of the trial: *"This is a study, so there may be problems"*. In a few cases, the risks were minimized *"This will not happen to me, I never had trouble with other methods"*.

*Final decision:*

Women arrived to a final decision after processing the information about the methods, the trials and the clinic. This decision stemmed from the evaluation they made of perceived benefits and risks, their rights and responsibilities, and their ideas of what research and volunteering in a trial mean.

Women highly valued having plenty of time to make the decision, and the confidence that it would be respected:

*“...They give you the leaflet, you read it calmly... You come to the counseling session, with other people, and they explain to you, they discuss the issues... You have plenty of time to make your decision, and it is my decision, no one else is going to make it...”*

When a woman was interested and received support from her husband or mother, she decided to participate. On the contrary, the two women who decided to withdraw from the trial did so because of pressure from partners who had a negative opinion of the method.

Side effects and risks and the balance between them and personal goals, were the main reasons to decide not to participate:

*“I never wanted to take the pill because of the weight gain, or other side effects. When I knew that those implants had hormones, I thought this was a negative trait”.*

*“Being a study, pregnancy prevention was not 100% assured, then I decided not to participate, I didn’t want to take the risk, since I had had two miscarriages”.*

Women who had been in the trial for six or more months said they would make the same decision again. These women, and also those who had discontinued method use, said that the final balance regarding their participation was positive, and many of them found no negative aspects. All of them agreed that, given the ability to pay for a method, they would prefer to participate in a trial again rather than obtaining it at a private service:

*“I want to continue, I would participate in other studies, I would bring friends so that they receive the same care I received.”*

*“In private practice they will just give me the method... no, no, I want to have all the information that I need and to be under close follow up.”*

*The meaning of signing the consent form:*

At the time women signed the IC form, most of them knew the requirements to participate, the general sequence and the procedures of the trial. However, some women did not handle complete information, ignoring, for example, how long did the trial last, or the precise sequence of the procedures.

Women frequently perceived that the consent form was useful both for the institution and the volunteers. It benefits the clinic because it proves that participants became involved in the study without external pressures and that they commit to comply with the study procedures. It benefits the volunteers by warranting their right to the free medical care associated to the trial.

*“I think (signing the consent form) means that I agree to what the document says. (I prefer) signing a document, because that makes you feel reassured, if something happens to you, they*

*have to hold themselves responsible, if you get sick they have to pay for the expenses”.*

Some women thought that the consent form is more useful for the institution, since it limits the clinic’s responsibility and, therefore, it is the volunteer who assumes the risks:

*“I think it is more useful for the institution, in case of risk or anything... the document said that in case of illness they would pay for all the medical costs, but they assumed no responsibility if the baby was born with any malformation”.*

Only one woman explicitly associated the consent form to the commitment of the institution to comply with all the procedures and perform all the exams described in the document.

*Perceived rights and responsibilities:*

All women knew they could withdraw from the study and have the method removed at any time. Nevertheless, they felt committed to stay in the trial up to the end: *“I knew I could withdraw at any moment, but I like to finish what I start”.* All were aware of their right to cost-free

medical care in case of health events related to the trial. A woman who thought that the clinic would provide care for pregnancy and delivery in case of method failure misunderstood this. The right to confidentiality was mentioned only once.

*Problems detected:*

There was occasional misunderstanding of the information given, either verbally or in the consent form. Some women had misconceptions, such as thinking there was a high risk of pregnancy during the trial. One participant understood that, if pregnancy occurred, the clinic would give her free medical care.

Some women commented about various aspects of the consent process at this clinic that could be improved. They asked for more information about reproductive health, and to involve their partners in the counseling process. They requested a quiet place at the clinic to read the consent form, rather than the waiting room. They suggested that each woman should be cared for by only one midwife, and emphasized that the midwife should ask women more about their doubts.

## **DISCUSSION**

The evaluation of the informed consent procedures described identified several essential components of a free decision-making process. Aspects highly valued by the participants were to be fully informed and to receive specific counseling, to have the opportunity to discuss the information with close parties and to have enough time to think their decision. Being informed, as well as the quality of care provided, contributed to build confidence in the research team.

Nevertheless, not all the information was clearly understood by all the participants, which may be partially explained by the little exposure to this type of information in the past or by previous misconceptions and myths. Women did not perceived some aspects of the research protocol as requirements of the study or as mutual benefits, but as "goods" that the institution offered them. Sometimes they interpreted the information about health care beyond what was actually offered. This was based on ambiguities in oral or written explanations, as well as their own needs and expectations.

Women made their personal risk-benefit balance. Meeting contraceptive needs, the quality of ICMER's services, and attributing a positive meaning to research, were usually placed in one side of the scale, and



side effects or risks in the other. Those who decided to participate had handled their doubts or fears through various mechanisms. Among those worried about contraceptive effectiveness, some carefully evaluated the risk of pregnancy, some considered that other benefits associated with the trial made the risk worthy enough, and some just thought they would be lucky enough not to get pregnant.

For some women, participation meant a reinforcement of self-esteem, the expression of altruism and a means to satisfy the need to transcend to do something “important” or that benefit others beyond their close environment. Even when women did not mention it explicitly, and in spite of the problems detected, the IC procedure described seem to support women’s autonomy. When women had good relationships, the process favored communication with the partner, and his participation in reproductive decisions.

However, we identified some factors that might limit women’s free choice, most of them socio-cultural and gender issues that are a matter of concern. These are reflected in women’s low awareness of their rights; their scarce questioning of the risks involved in the clinical trial or the intentions and authority of providers; the difficulty that some had to

express fears or doubts during their visits to the clinic; and the fact that they need a signed document to commit to their own health care.

Sometimes, women expressed questions or concerns about the procedures as opinions of the partners. This might be an expression of gender differences: men have cultural permission to be suspicious, while women's socialization includes being humble and acceptant. Another interpretation is that, in our culture, men are not in charge of pregnancy prevention. Thus, they maintain objectivity with regard to the risks involved in a trial, while the women tend to minimize them because of their strong motivation for contraception.

The evaluation of benefits and risks could be jeopardized because of women's needs. Good reproductive health care may be a benefit only when it is not available. The economic motivation might be important for disadvantages groups, since cost-free contraceptives clinical follow-up, and paying for transportation were attractive to the participants.

The desire to benefit others could be also interpreted as a gender role, which reinforce in women the will to take care of other people. Participants in this study belong to a segment of the population that lacks other opportunities to express altruism. If their conditions change

and they may express their altruistic tendencies in other ways, less women may be interested in participating in clinical studies.

The decision to enter the trial is the peak of an emotional process where the information received and assimilated meets the needs, expectations, and wishes of the women. Signing the consent form is an explicit means to communicate this intimate decision to others, and make a commitment. The drawback of this is that, although aware of the right to withdraw from the study, might feel committed to continue out of sense of duty.

The findings of this study suggest some recommendations for researchers in developing countries, being the most important that researchers should be aware that participants might not perceive themselves as subjects of rights. Thus, special efforts must be made to promote their awareness in this matter.

The information given to potential and actual volunteers should clearly state that the procedures of the study are essential for the research. Volunteers collaborate, and the institution needs them. If the research conditions fit their needs and preferences, they can freely accept and benefit from them. This represents an exchange of mutual benefits between participants and the research institution. The research

institution needs to make sure that the participants understood the obligations acquired by each part, and what the institution is not committed to do for them.

A major challenge for researchers is to deliver complex information at the right time in appropriate language and style. Therefore, the IC document should be adapted to the local culture and the educational level of the population. This might become a hard task, because of the technical and ethical issues to be conveyed. Thus, developing a comprehensive document may take several reviews and assessment of volunteers' understanding. In turn, when subjects clearly understand this information, they make better decisions, comply with research procedures, experience less surprise later on, and stay longer in the trials. Therefore, researchers benefit from enrolling informed volunteers. IC includes decisions during the trial, especially when women experience problems or nuances. Therefore, IC is a continuous process that requires continued counseling, problem solving, complementary services, and integrating the partner when necessary. In addition, the researchers should make sure that the volunteers are fully aware of their right to withdraw from the trial, and are willing to do so if necessary.

The participant's rights, ethical and legal obligations, and the interest of researchers do not contradict one another. The IC process is useful for the participant, and allows her/him to take an active role in self-care and bring relevant information to the research process. In turn, the research benefits from promoting active participation and self-care by augmenting the probability of obtaining timely, accurate and complete information about volunteer's experience.

All the factors involved in the decision-making process described may enhance participant's ability to exercise their autonomy and their right to choose. They may also assure research institutions that they are conducting the trials without exercising any type of coercion over the participants. Therefore, it is relevant for every research setting to design strategies to ensure the confluence of volunteers' and researchers' interests in a harmonic and equitable way.

### **Acknowledgements**

The authors are grateful to the participants and the clinic staff for their collaboration. The study was funded by the Gender Panel of the UNDP / UNFPA / WHO / World Bank Special Program of Research, Development and Research Training in Human Reproduction.

## REFERENCES

1. Nuffield Council on Bioethics. 1999. *The Ethics of Clinical Research in Developing Countries: A discussion paper*. London, UK.
2. Council for International Organizations of Medical Sciences, World Health Organization. 1993. *International Ethical Guidelines for Biomedical Research involving Human Subjects*. Geneva.
3. World Health Organization. 1995. *Guidelines for good clinical practice (GCP) for trials on pharmaceutical products*. WHO Technical Report Series 850:97-137, Geneva.
4. Pellegrini Filho and Macklin R, editors. 1999. *Research on Human Subjects: International Experience*. Regional Program on Bioethics PAHO/WHO. Santiago, Chile.
5. American College of Obstetricians and Gynecologists. 1993. *Ethical Dimensions of Informed Consent*. Women's Health Issues, The Jacobs Institute of Women's Health, 3(1) 1-10.

6. G Salazar, M Tijero and S Sánchez. 1993. Decisions and Perspectives of Women who participate in Contraceptive Clinical trials. Final Report to the UNDP / UNFPA / WHO / World Bank Special Program of Research, Development and Research Training in Human Reproduction.
7. H Vera. 1993. The client's view of high-quality care in Santiago, Chile. *Stud Fam Plann* 1993; 24:40-49.
8. J Taylor and R Bogdan. 1992. *Introduction to Qualitative Research Methods. The Search for Meaning*. New York. J Wiley and Sons Inc.